

# PATHOLOGY OF BLOOD AND LYMPHATIC SYSTEM

Dr. Tariq Al-Adaily, MD

Associate Professor


Department of Pathology

The University of Jordan


Email: [TNALADILY@ju.edu.jo](mailto:TNALADILY@ju.edu.jo)



School of Medicine



- First Semester 2021 / 2022
- Reference: Robbins Basic Pathology 10<sup>th</sup> ed



# ANEMIA



# DEFINITION

- ❑ Reduction of oxygen carrying capacity of blood secondary to decrease in red cell mass
- ❑ Leads to tissue hypoxia
- ❑ Practically, measure by Hemoglobin concentration, and Hematocrit

د. طارق العديلي




Faculty of Medicine



# ANEMIA AND ERYTHROPOIETIN

- Anemia triggers production of erythropoietin
- Causes compensatory erythroid hyperplasia in bone marrow (BM)
- In acute anemia, production can increase by 5x or more in healthy people
- In severe cases, causes extramedullary hematopoiesis in secondary hematopoietic organs (spleen, liver and lymph nodes)
- Exceptions: anemia of renal failure, anemia of chronic inflammation



# CLASSIFICATION ACCORDING TO CAUSE

## 1) Blood loss

## 2) Diminished RBC production

- Iron deficiency anemia
- Anemia of chronic inflammation
- Megaloblastic anemia
- Aplastic anemia
- Pure red cell aplasia
- Myelophthisic anemia
- Myelodysplastic syndrome
- Anemia of renal failure
- Anemia of hypothyroidism

## 3) Increased destruction



## (hemolytic anemia)

- Extrinsic factors (infection, antibody, mechanical)
- Intrinsic RBC abnormalities:
  - 1) Hereditary (membrane, enzyme, Hg abnormalities)
  - 2) Acquired (Paroxysmal nocturnal hematuria)





# CLASSIFICATION ACCORDING TO MORPHOLOGY BLOOD FILM

- Size: normo, micro, macrocytic (MCV)
- Color: normo, hypochromic (MCH)
- Shape: anisopoikilocytosis (spherocytes, sickle, schistocytes) (RBC distribution width)
- Hypochromic microcytic anemia usually reflects impaired Hg synthesis
- Macrocytic anemia reflects stem cell disease and maturation



# RBC INDICES

- Can be directly measured, or automated
- Slight variation is present between labs, geographic areas
- Sex, age, race, mobility status have effect
- Reticulocyte count: helps differentiate hemolytic anemia (high) from aregenerative anemia (low)



	Units	Men	Women
Hemoglobin (Hb)	g/dL	13.2–16.7	11.9–15.0
Hematocrit (Hct)	%	38–48	35–44
Red cell count	$\times 10^6/\mu\text{L}$	4.2–5.6	3.8–5.0
Reticulocyte count	%	0.5–1.5	0.5–1.5
Mean cell volume (MCV)	fL	81–97	81–97
Mean cell Hb (MCH)	pg	28–34	28–34
Mean cell Hb concentration (MCHC)	g/dL	33–35	33–35
Red cell distribution width (RDW)		11.5–14.8	

\*Reference ranges vary among laboratories. The reference ranges for the laboratory providing the result should always be used in interpreting a laboratory test.





# CLINICAL FEATURES OF ANEMIA

- Dizziness
- Fatigue
- Pallor
- Headache

Adaptive changes:

- Tachycardia
- Tachypnea
- Increased red cell 2,3-diphosphoglycerate

If the patient has heart or lung diseases, symptoms will be worse

د. طارق العديلي



# CLINICAL SYMPTOMS IN SPECIAL TYPES OF ANEMIA

- Chronic hemolytic anemia: jaundice, pigmented gall bladder stones, redurine





Faculty of Medicine



# CLINICAL SYMPTOMS IN SPECIAL TYPES OF ANEMIA

- Extramedullary hematopoiesis: splenomegaly, hepatomegaly
- Thalassemia major and sickle cell anemia: growth retardation, bone deformity, secondary hemochromatosis (damage to heart, endocrine glands)



# ANEMIA OF ACUTE BLOOD LOSS

- Symptoms are related to decreased intravascular volume,
- If loss is > 20% of blood volume, patient might have hypovolemic shock and death
- Body responds by shifting fluid from interstitial to intravascular space, causing dilutional anemia and worse hypoxia (stays 2-3 days)
- Erythropoietin secretion is stimulated, activating BM erythropoiesis (needs 5-7 days)
- In internal hemorrhage, iron is restored from extravasated RBCs and used again in erythropoiesis
- In external and GIT hemorrhage, iron is lost, which complicates anemia
- The anemia is normochromic normocytic, with reticulocytosis



# ANEMIA OF CHRONIC BLOOD LOSS

- ❑ Occurs when the rate of RBC loss exceeds regeneration
- ❑ Mostly occurs in gastrointestinal diseases, also in excessive menstruation
- ❑ Results in iron deficiency, anemia appears hypochromic and microcytic, low reticulocytes



Faculty of Medicine

